

New primate genus from the Miocene of Argentina

Marcelo F. Tejedor^{*†}, Adán A. Tauber[‡], Alfred L. Rosenberger^{§¶}, Carl C. Swisher III^{||}, and María E. Palacios^{**}

^{*}Consejo Nacional de Investigaciones Científicas y Técnicas, Laboratorio de Investigaciones en Evolución y Biodiversidad, Facultad de Ciencias Naturales, Sede Esquel, Universidad Nacional de la Patagonia "San Juan Bosco," Sarmiento 849, 9200 Esquel, Argentina; [†]Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Avenida Velez Sarsfield 249, 5000 Córdoba, Argentina; [‡]Department of Archaeology and Anthropology, Brooklyn College, City University of New York, 2900 Bedford Avenue, Brooklyn, NY 11210; [§]Division of Vertebrate Zoology (Mammalogy), American Museum of Natural History, Central Park West at 79th Street, New York, NY 11024; [¶]Department of Geological Sciences, Rutgers University, Piscataway, NJ 08854; and ^{**}Museo Regional Provincial "Padre Manuel Jesús Molina," Ramón y Cajal 51, 9400 Río Gallegos, Argentina

Edited by Jeremy A. Sabloff, University of Pennsylvania Museum of Archaeology and Anthropology, Philadelphia, PA, and approved December 28, 2005 (received for review August 4, 2005)

***Killikaike blakei* is a new genus and species of anthropoid from the late Early Miocene of southeastern Argentina based on the most pristine fossil platyrrhine skull and dentition known so far. It is part of the New World platyrrhine clade (Family Cebidae; Subfamily Cebinae) including modern squirrel (*Saimiri*) and capuchin monkeys (*Cebus*) and their fossil relatives known from Early to Middle Miocene and subrecent periods. Living cebines are relatively large-brained, adroit predatory foragers and live within complex social groups, and wild capuchins exhibit a wide range of behaviors associated with enhanced intelligence. We show that *K. blakei* lacks diagnostic derived characteristics of the lower face and premolar dentition that are shared by modern cebines, but its strongly vaulted frontal bone and capacious anterior cranial fossa indicate the early evolution of an enlarged forebrain.**

Miocene | Paleoprimatology | Patagonia | Platyrrhini | Cebinae

Intact craniodental remains of pre-Pleistocene fossil New World anthropoids are exceedingly rare (1–3). Of the five Miocene genera that are known from crania (*Dolichocebus*, *Tremacebus*, *Homunculus*, *Chilecebus*, and *Lagonimico*), the first three are essentially edentulous and all but one are broken and/or deformed. *Chilecebus carracoensis* from a site in the western Andes of Chile (4) is an exception. It preserves an unspoiled skull with teeth *in situ*, albeit heavily worn. Here we report a fossil of a platyrrhine that preserves the entire face and also an unworn, little damaged dentition that is the best anatomical evidence of the maxillary teeth of any Santacrucian fossil monkey. It also provides relatively complete and undistorted evidence of the anterior braincase of a fossil platyrrhine. The two specimens of this species come from the classic Santacrucian sediments of Patagonian southeastern Argentina (5, 6), which has produced a substantial primate fauna (1, 7, 8, ^{††}). They are the southernmost fossil platyrrhine primates ever described.

Systematic Paleontology. The classification of *Killikaike blakei* is as follows: Order Primates (Linnaeus, 1758); Suborder Anthropoidea (Mivart, 1864); Superfamily Ateloidea (Gray, 1825); Family Cebidae (Bonaparte, 1831); Subfamily Cebinae (Bonaparte, 1831; Mivart, 1865); *K. blakei*, gen. et sp. nov.

Holotype. MPM-PV 5000 (Museo Regional Provincial "Padre Manuel Jesús Molina" Padre Molina, Vertebrate Paleontology Collection) is an intact, undistorted face of an adult individual that preserves the forehead, orbits, snout, dental arcade, and roots of the pterygoid processes. The dentition includes right C, P^{2–3}, root of P⁴, crowns of M^{1–3}; left C, broken crowns of P^{2–3}, and roots of M^{1–3} (Fig. 1A–E).

Hypodigm. Holotype and MPM-PV 1607, a right M^{1–3} associated in a piece of maxilla (Fig. 1F).

Locality and age. MPM-PV 5000 was collected >25 years ago at the locality Killik Aike Norte (formerly known as Estancia Felton) (5, 6, 9), ≈50 km northwest of the city of Río Gallegos, Santa Cruz Province, Argentina (Fig. 2). MPM-PV 1607 was collected in January of 2005 at the same locality. Killik Aike Norte is part of the Santa Cruz Formation (10) exposed in the northern cliffs

of the estuary of the Río Gallegos river. The stratigraphy is divided into two members (6): Estancia La Costa (120 m thick with 18 fossiliferous levels) and Estancia Angelina (103 m thick with four fossiliferous levels). There are two types of volcanic ash at the locality. The exact provenience of the MPM-PV 5000 specimen is not known, but the mineralogy and chemistry of the rock that encased it are consistent with an origin in the lower volcanic tuff, which is part of the Estancia La Costa member (Fig. 3). MPM-PV 1607 was also collected from the lower tuff. The level 6 tuff of the Estancia La Costa member at the locality Estancia La Costa, northeast of Killik Aike Norte, previously yielded a primate skull attributed to *Homunculus patagonicus* (7). Correlation of the level 6 at La Costa with the tuff at Killik Aike Norte is not confirmed yet.

We have determined the age of the MPM-PV 5000 through ⁴⁰Ar/³⁹Ar laser incremental-heating analyses of plagioclase separated from the tuff level where the skull was found at Killik Aike Norte. Three separate incremental-heating experiments yielded reproducible plateau ages of 16.45 ± 0.14, 16.5 ± 0.2, and 16.5 ± 0.3 mega-annum (Ma) (see Fig. 4, which is published as supporting information on the PNAS web site).

The Ca/K ratio, derived from the ³⁷Ar/³⁹Ar ratio obtained during the ⁴⁰Ar/³⁹Ar dating of the plagioclase, can be used to delimit possible correlative Santa Cruz tuffs previously dated at the coastal sites of Monte Observación and Monte León, which are ≈100 km north of Killik Aike Norte. These sites contain key fossil mammal localities of the Santa Cruz Formation and have yielded ⁴⁰Ar/³⁹Ar ages similar to that reported here for Killik Aike Norte. The low Ca/K ratios of the Killik Aike Norte plagioclase compare best with Monte Observación MO2, although the variation in the Ca/K measurements does not preclude possible correlation with MO64 or ML18 tuffs, and further geochemical study is needed (11, ^{‡‡}).

Our ⁴⁰Ar/³⁹Ar age for Killik Aike Norte is appreciably younger than the older age limit of 17.7 Ma for the Santa Cruz Formation based on a K-Ar date from the same site (12). The older age may be attributed to contaminant (detrital grains incorporated into the bulk K-Ar sample). The younger age for Killik Aike Norte reported here corroborates the assessment of Fleagle *et al.* (10, 11) that the base of the Santacrucian at the classic coastal localities between Río Gallegos and Río Coyle is not older than ≈16.5 Ma. The older age limit for the Santacrucian based on the 17.7 Ma K-Ar date may be in error. The main terrestrial fossiliferous units of the Santacrucian Land Mammal Age appear to have been laid down over a relatively short time

Conflict of interest statement: No conflicts declared.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviation: Ma, mega-annum.

[†]To whom correspondence should be addressed. E-mail: mtejedor@lieb.org.ar.

^{††}Fleagle, J. G., Buckley, G. A., & Schloeder, M. E. (1988) *J. Vert. Paleontol.* 8, 14a (abstr.).

^{‡‡}Fleagle, J. G., Perkins, M., Bown, T. M., Tauber, A. A., & Dozo, M. T. (2004) *J. Vert. Paleontol.* 24, Suppl. 3, 58A (abstr.).

© 2006 by The National Academy of Sciences of the USA




Fig. 1. Holotype cranium and referred upper molars of *K. blakei*, a new genus of platyrrhine primate from the Miocene of Argentina. Frontal (A), right lateral (B), palatal (C), and posterior (D) views of MPM-PV 5000, and an illustration of the cheek teeth (E). (F) MPM-PV 1607 in occlusal view. (Scale bar, 1 cm.)

period of $\approx 400,000$ years, approximately between 16.1 and 16.5 Ma.

Etymology. *Killikaike*, after Killik Aike Norte, the name of the ranch and fossil site; *blakei*, in honor of the Blake family, who donated the fossil to the Padre Molina museum.

Diagnosis. *Killikaike* is unique among platyrrhines in the following combination of craniodental characters: relatively short face that is not wide anteriorly, narrow interorbital region, deep recessed orbits that lack an imperforate medial orbital wall, high arched frontal bone conforming to an enlarged anterior cranial fossa, and nonbunodont cheek teeth with subtriangular premolars and hypocone-bearing quadrate molars. It is a platyrrhine similar in size to *Saimiri*, *Callicebus*, or *Leontopithecus* based on dental and cranial dimensions (Tables 1 and 2), with an upper tooth count of 2.1.3.3. Unlike *Cebus* and *Saimiri*, the face is relatively narrow and the snout tapers anteriorly, making it shorter than *Aotus* and *Pithecia*, for example, but without abbreviated premaxillae as in *Callicebus*. The interorbital region is relatively narrow, in contrast to callitrichines, pitheciines, atelines, and *Chilecebus*, but less so than in existing cebines and *Dolichocebus*. Orbits are not enlarged as in *Aotus* and *Tremacebus*, and the medial orbital walls are closely spaced and not fenestrated, in contrast to *Dolichocebus* and *Saimiri*. The orbits are recessed beneath the

arched and uptilted frontal bone, as in modern cebines and unlike most other platyrrhines, where the frontal trigon is flattened in the midline. A relatively large lateral orbital foramen is partially preserved, unlike most other platyrrhines. Postcanine crowns are moderately crested, resembling *Saimiri* and unlike the bunodont *Cebus*. However, in contrast to *Saimiri* and *Cebus*, where P^{2-3} are transversely wide and lingually expanded and sometimes have a lingual cingulum, in *Killikaike* these teeth are relatively narrower and more triangular, and they lack lingual cingulum. M^{1-2} are quadrate with high cusps and well developed, offset hypocones, strong lingual cingula, and well developed crests, distinguishing the genus from all pitheciines, atelines, callitrichines, and *Szalatavus*. The molars are relatively narrower than in *Branisella* and *Chilecebus*. M^1 is considerably wider than M^2 , and M^3 is reduced but otherwise an unremarkable two-cusped crown.

Results and Discussion

MPM-PV 5000 is the forehead, face, and palate of a young adult, probably female. It evidently broke away from the neurocranium, which has not been recovered, behind the craniofacial junction. The snout and orbits are remarkably well preserved, bilaterally undamaged except for a slight crack at the left lateral

Table 1. Dental measurements (means in millimeters) of fossils and selected living species

	C		P ²		P ³		M ¹		M ²		M ³	
	BL	MD	BL	MD	BL	MD	BL	MD	BL	MD	BL	MD
MPM-PV 5000	3.4	2.8	3.6	2.8	4	3	—	4.1	4.9	3.7	4.5	2.9
MPM-PV 1607	—	—	—	—	—	—	5.7	4.1	5.5	4	4.9	3.4
<i>Saimiri sciureus</i> (3m, 3f)	2.9	3.1	3.6	2.4	3.9	2.2	4	3	3.6	2.6	2.7	1.7
<i>Saimiri oerstedii</i> (3m, 3f)	3	3.3	3.2	2.3	3.5	2	3.6	2.6	3.4	2.4	2.4	1.6
<i>Cebus apella</i> (4m, 4f)	6	6.7	6.5	3.9	3.5	3.5	6	4.4	5.4	3.9	4.4	3
<i>Callicebus moloch</i> (4m, 4f)	3.1	2.6	3.2	2.1	3.6	2.2	4.4	3.2	4.3	3	3.3	2.3
<i>Aotus vociferans</i> (3m, 3f)	2.5	2.8	2.9	1.9	3.2	2	3.9	3.1	3.7	2.9	3.2	2.1
<i>Leontopithecus rosalia</i> (2m, 1f, 3?)	3	3.3	3.2	2.6	3.8	2.3	4.1	3.2	3.3	2.4	na	na

Sexes and sample sizes are in parentheses. Data on body weight and cranial capacity and the Index of Cranial Capacity are from Martin (17) and are not based on the same individuals. BL, buccolingual breadth; MD, mesiodistal length; m, males; f, females; na, not applicable.

forms, which represent nearly one-quarter of the pre-Pleistocene fossil record of platyrrhines, has been fueled by the poor quality of the fossils (1–3, 21, 22, §§). *K. blakei*, as the best-preserved cranium of a Tertiary New World monkey, thus has important bearing on the evolution of the platyrrhine radiation.

The narrow interorbital region, elongate descending process of the frontals, rounded, recessed orbits, and vaulted frontal bones are a combination of features found only among extant platyrrhines in *Saimiri* and *Cebus* and in the fossil *Dolichocebus*. These features together provide strong evidence that *K. blakei* is a member of the cebine clade, whose monophyly is well corroborated by morphology and molecules (22–24). The origins of this clade may have occurred >22 million years ago (24). Absent in *Killikake* are derived rostral and dental features shared by these sister taxa, such as the U-shaped dental arcade, large transversely broadened and rectangular upper premolars, a foreshortened face, and enlarged sexually dimorphic canines (16, 25).

On the other hand, the domed frontal bone and measured estimates of anterior cranial fossa size given above leave little doubt that the forebrain of *K. blakei* was shaped and proportioned like *Saimiri*. *Saimiri* and *Cebus* are two highly encephalized anthropoids (15, 25). Martin (15) calculated an Index of Cranial Capacity for primates that measures the amount of encephalization in a species above what is expected for its body size, based on a “basal” insectivore regression model (Table 2; also see Fig. 5, which is published as supporting information on the PNAS web site). The three highest index values were as follows: *Homo sapiens*, 23; *Australopithecus africanus*, 12.4; and

Cebus sp., 11.7. Capuchins ranked well above chimpanzees (8.2) and orangutans (7.7). The *Saimiri* value was 7.0, well within the range of apes and Old World monkeys. Thus, as with *Cebus* and *Saimiri*, expansion of the forebrain in *K. blakei* is a likely indication that overall brain size was also large relative to body size.

At ≈16.4 million years, *K. blakei* presents a primitive cebine facial and dental structure but an advanced brain, whereas contemporaneous apes had brain sizes that were unexceptional and probably smaller than the modern norms (26). The importance of facial foreshortening and broadening of the snout and premolars in modern cebines remains to be determined, but these features are probably connected with processing tough insects and ripping or tearing fibrous woody materials to harvest embedded food. The absence of these features in *Killikake* suggests a brain-first/teeth-second adaptive sequence in the mosaic evolution of their extractive foraging–feeding specialization, with problem-solving skills and manual dexterity preceding oral-harvesting and food-processing adaptations.

Materials and Methods

We have compared the fossil platyrrhines described here with the material of living and fossil platyrrhines deposited in the collections of the Museo Argentino de Ciencias Naturales (Buenos Aires) and the American Museum of Natural History (New York).

Table 2. Cranial measurements (means in millimeters) of fossils and selected living species

	Interorbital breadth	Biorbital breadth	Orbital height	Orbital width	Postorbital constriction	Bicanine breadth	Bimolar width	Body weight, g	Cranial capacity, cm ³	Index of cranial capacity
MPM-PV 5000	4.4	30.5	13.9	13.1	27.3	14.4	23.9	—	—	—
<i>Saimiri sciureus</i> (3m, 3f)	3.2	33	15.3	13.9	29.8	17.3	20.4	914	23.6	7
<i>Saimiri oerstedii</i> (3m, 3f)	3	32.5	14.7	13.9	29.7	15.9	19.6	—	—	—
<i>Cebus apella</i> (4m, 4f)	5.2	51.2	21.3	20.2	38.6	28	28.2	2537	76.2	11.7
<i>Callicebus moloch</i> (4m, 4f)	6.7	36	15.4	14.1	31.4	14.3	21.3	1078	18.3	4.9
<i>Aotus vociferans</i> (3m, 3f)	4.9	42	19.3	19.1	31	15.4	20.4	985	16.9	4.8
<i>Leontopithecus rosalia</i> (2m, 1f, 3?)	6.4	28.6	11.7	11.3	23.2	15.9	20.7	—	—	—

Sexes and sample sizes are in parentheses. Data on body weight and cranial capacity and the Index of Cranial Capacity are from Martin (17) and are not based on the same individuals. m, males; f, females.

